

UNCLASSIFIED

AD NUMBER	
AD318487	
CLASSIFICATION CHANGES	
TO:	unclassified
FROM:	confidential
LIMITATION CHANGES	
TO:	Approved for public release, distribution unlimited
FROM:	Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; Jun 1960. Other requests shall be referred to U.S. Army Chemical Warfare Lab., Army Chemical Center, MD.
AUTHORITY	
USAEA Notice, 9 Jul 1971; USAEA Notice, 9 Jul 1971	

THIS PAGE IS UNCLASSIFIED

UNCLASSIFIED

AD_ 318487

DEFENSE DOCUMENTATION CENTER

FOR

SCIENTIFIC AND TECHNICAL INFORMATION

CAMERON STATION ALEXANDRIA, VIRGINIA

CLASSIFICATION CHANGED
TO UNCLASSIFIED
FROM CONFIDENTIAL
PER AUTHORITY LISTED IN

DD FORM 1-71

15 September 1971



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

318 487

CONFIDENTIAL

U. S. Army
Chemical Warfare Laboratories
Technical Report

CWLR 2399

Breathing Patterns During Human Exposure to CS (U)

by

F. N. Craig

W. V. Blevins

E. G. Cummings

June 1960



ARMY CHEMICAL CENTER, MD.

CONFIDENTIAL

CONFIDENTIAL

June 1960

CWLR 2399

BREATHING PATTERNS DURING HUMAN EXPOSURE TO GS (U)

by

F. N. Craig
W. V. Blevins
E. G. Cummings

Physiology Division

Recommending Approval



DOUGLAS LINDSEY
Lt. Col., MC
Director of Medical Research

Approved:



S. D. SILVER
Deputy Commander for
Scientific Activities

U. S. ARMY
Chemical Corps Research and Development Command
CHEMICAL WARFARE LABORATORIES
Army Chemical Center, Maryland

CONFIDENTIAL

UNCLASSIFIED

(U)

FOREWORD

These observations were authorized under Task 4C08-02-023-01, Basic and Applied Physiology (U). They were made between 30 March and 13 April 1959. This report was submitted for publication in March 1960.

Acknowledgement

The authors are indebted to the staff of the Aerosol Branch for making available to us the human volunteer subjects during exposures conducted by the Aerosol Branch, and for their observations concerning the exposures.

Notices

This document contains information affecting the national defense of the United States within the meaning of the Espionage Laws, Title 18, U. S. C., sections 793 and 794. The transmission or the revelation of its contents in any manner to an unauthorized person is prohibited by law.

Reproduction of this document in whole or part is prohibited except with permission of the issuing office; however, ASTIA is authorized to reproduce the document for U. S. Governmental purposes.

Disposition

When this document has served its purpose, DESTROY in accordance with paragraph 35 of AR 380-5. DO NOT return to U. S. Army Chemical Warfare Laboratories.

UNCLASSIFIED

CONFIDENTIAL

(C)

DIGEST

During the BLACK MAGIC program of tests on the incapacitating agent CS, detailed observations of the breathing behavior of six human volunteer subjects were made by means of the remotely-controlled breath-recording system under development for the CARAMU program. Although the breathing patterns were disrupted by CS, adequate ventilation of the lungs was maintained, so that the incapacitation is attributed to the unpleasant sensations rather than to any degree of respiratory failure.

CONFIDENTIAL

UNCLASSIFIED

(U)

CONTENTS

	<u>Page</u>
I. INTRODUCTION.....	5
II. PROCEDURE	5
III. RESULTS	7
IV. DISCUSSION.....	9
V. SUMMARY	11
VI. CONCLUSION	12
VII. LITERATURE CITED	12
APPENDIX, Tables and Graphs	15

UNCLASSIFIED

CONFIDENTIAL

BREATHING PATTERNS DURING HUMAN EXPOSURE TO CS (U)

I. (U) INTRODUCTION.

During the spring of 1959 a large number of human volunteers were exposed to the riot control agent, CS, in the form of an aerosol generated under controlled conditions in a wind tunnel. The observers¹ concluded that the agent produced incapacitation by affecting the eyes, the respiratory tract, or both. They regard the symptoms referable to the respiratory tract as potentially the most capable of causing incapacitation. Accordingly, it became of interest to observe the breathing behavior. The availability of one unit of a self-contained, remotely-controlled breath-recording system, under development for estimating casualties from munitions (CARAMU), permitted us to make observations in a short series of exposures.

II. (U) PROCEDURE.

Of the four or five men entering the wind tunnel for an exposure, one man was equipped with the breath-recording device. The recorder was started by a radio signal and when an outside observer saw the cloud arrive he impressed by radio a signal mark on the record. To measure the duration of the exposure, a stop watch was started at the same time as the radio signal and stopped when the man left the wind tunnel. The concentration of CS was determined by members of the Aerosol Branch.

The device consists of (a) an air-flow transducer mounted in an oronasal mask of rubber sheet 0.020 inch thick attached to the skin with adhesive tape, (b) a recorder (figure 1, appendix) with enough magnetic tape to last 3 minutes and (c) a radio receiver and other electronic components. Items b and c were carried in a pack on the back (figure 2, appendix) with a shielded cable leading to the transducer. The equipment carried by the man weighed 10 pounds. A developmental prototype of the transducer has been described by Dahlke and Welkowitz.^{2,3} The finished transducer is 2 inches in diameter and 1/2 inch thick and is illustrated elsewhere.⁴ It consists of networks of strain-gage wire bonded to alternate sides of 8 flexible vanes attached to the circumference of the airway and projecting into the airway. When no air is moving, the airway is nearly closed; when air moves in either direction, the vanes bend, opening the airway and deforming the strain-gage wire. Since the gages form two arms of a bridge circuit, air movement produces an imbalance in the bridge. The arrangement of the gages compensates for

CONFIDENTIAL

UNCLASSIFIED

thermoelectric effects and the connections are waterproofed to protect against condensation from the expired air. The output is nearly the same in either direction if the direction of air movement alternates as in breathing. The balance changes, however, if the transducer is subjected for very long to steady or intermittent flow in only one direction.

The output of the bridge enters a phase modulation circuit and is then subjected to pulse position modulation for recording on magnetic tape. The circuit and recorder are modifications by Dahlke and Welkowitz² of those described by Upham and Dranetz.⁵ The tape transport mechanism is driven by a voltage-sensitive motor powered by a nickel-cadmium rechargeable battery. Successful operation depends on a delicate balance between the battery power and the resistance of the tape transport mechanism. Roughness in the tape transport or imbalance in the demodulator results in noise in the baseline of the final record. A decline in the battery voltage during a 3-minute run produces a shift in the baseline level and reduction in amplitude of the signal. For retrieval of information, the tape is played back. The output from the tape passes through a demodulator and is recorded on an oscillograph in the same manner as a conventional pneumotachogram. The remote control radio transmitter and receiver are of the type used with model airplanes.³ The system is identified as the production prototype delivered by the contractor in December 1958.⁶

As shown in figure 3 (appendix), the resistance of the transducer to air movement was comparable to the inspiratory resistance of the M17 protective mask.⁷ The transducer used in these experiments had a nearly linear calibration for steady air flows measured by Fisher-Porter rotameters (figure 3). The response of the transducer apart from the remainder of the tape recording system has been described elsewhere.⁸ When the same breath was passed through the conventional screen pneumotachometer and the transducer in series, the response was comparable with respect to ability to follow sudden changes in air flow and in the tidal volume of the breath. The transducer is too thin, however, to permit sufficient bending of the vanes to respond to instantaneous flows above 250 liters per minute.

The accuracy of the system was tested by comparing the volumes of several breaths during and after a brief run on the treadmill by a man wearing the transducer under an M9A1 protective mask. The canister was removed and the inlet was connected to a 120-liter recording spirometer with a scale factor of 0.1326 liter per millimeter of height of the bell. The transducer output was recorded on magnetic tape. In order to resolve the oscillations imposed on the record of instantaneous flow by the impact

UNCLASSIFIED

UNCLASSIFIED

of the feet on the treadmill and the consequent shock to the column of air extending from the lungs out through the mask, a high speed of paper transport in the Sanborn oscillograph was required. This amounted to 266 centimeters per minute. Since the playback speed is double the recording speed, the Sanborn paper speed was equal to 133 centimeters per minute of breathing. The duration of the breaths as recorded on tape was verified by comparison with a simultaneous recording of mask pressure on another Sanborn oscillograph. The volumes of several breaths were measured by reading from the record the instantaneous flow for each millimeter of the time scale. The instantaneous flow is the product of the pen deflection and the calibration factor for the system, 10 liters per minute per millimeter of pen deflection. The sum of the instantaneous flows divided by 1330 millimeters per minute yields the volume of the breath in liters. In table 1 (appendix) the inspired volumes recorded by the spirometer and by the transducer are listed together with the volumes of the intervening expirations also recorded by the transducer.

Since so large an excess of inhalation over exhalation is unlikely, sources of error were considered. In order to follow the rapid changes in flow, a hot pen was required; this produced a wide trace at slow flows and made a precise setting of the baseline uncertain. Inspection of the record shows the baseline may have been 0.5 millimeter further on the inspired side than the line employed in reading the pen deflections. From the total number of readings, 323 for inspiration and 297 for expiration, it was calculated that the mean pen deflections were 5.1 millimeters for inspiration and 4.5 millimeters for expiration. Corrected for possible error in the baseline of this size and direction, the average pen deflections become 4.6 millimeters for inspiration and 5.0 millimeters for expiration, and the total volumes 11.2 liters for inspiration and 11.2 liters for expiration. Such a shift in the average pen deflection is in harmony with the fact that the inspiratory phase was longer than the expiratory phase. The difference between 11.2 and the spirometric total in table 1 is not far from the 5 percent discrepancy between the spirometer and the rotameter used to calibrate the transducer. The volume collected in the spirometer is less by this much than the volume blown through the rotameter calculated from the rate indicated by the rotameter and the time of collection. This discussion does not exhaust the possibilities for error in the portable breath-recording system but may indicate that it has sufficient accuracy for field use.

III. (C) RESULTS.

(U) The records are illustrated in figures 4 and 6, appendix. A period of quiet breathing in the wind tunnel before the cloud arrived was

UNCLASSIFIED

CONFIDENTIAL

followed by the exposure, and the subsequent recovery outside the wind tunnel. Deviations from the quiet breathing pattern as a result of the exposure consist of periods of no air movement (a) between inspiration and expiration, (b) between expiration and inspiration, and (c) during the expiratory phase. Expanded portions of the records appear in figures 5 and 7, appendix. A spike in the expiratory direction was impressed on the record by a radio signal to indicate the arrival of the cloud. This time is only an approximation, since the concentration edge of the cloud is not sharp; some aerosol is thought to reach the subjects before the visible wisps of cloud can be seen. The following information applies to the top record in figure 5. The exhalation following the signal appeared normal and a further 0.43 liter was inhaled out of the total of 0.59 liter in that breath before the first irregularity in breathing could be seen. A complex cough followed and, after inhaling a total of 0.93 liter more, the subject left the wind tunnel at B. In one inspiration the instantaneous flow rose from 0 to 200 liters per minute within 0.05 second. The various periods in which no air flow was recorded add up to about 5.8 seconds. The subject did not intentionally hold the breath as a device to prolong the exposure, perhaps because of the risk of taking in a large breath at the end of the hold.

(C) Numerical data are listed in table 2, appendix. The concentration of CS in the aerosol cloud ranged from 5 to 150 micrograms per liter and the voluntary exposure times from 110 to 12 seconds. The total dose inhaled depended on the breathing during the exposure period so there was no close relation between concentration and dose. The changes in the breathing pattern as a result of exposure were the three types of interruption of breathing mentioned above plus the cough, but the quantitative aspect of the changes was quite varied from one exposure to another. The first detectable change in the breathing pattern came from 3 to 12 seconds after the arrival of the cloud, although in the exposure of BM no marked change in the breathing pattern was seen. In this case the subject had had considerable experience with the agent and entered the wind tunnel with the intention of maintaining a regular breathing pattern. The volume inhaled between the signal and the first respiratory response and the total inhaled during the exposure are listed to permit calculations of the doses inhaled. The times without breathing represent the sum of all the periods of zero flow observed during the exposure. In no case (except that of JC) was there a period long enough to suggest an intention to hold the breath. The total volume exhaled during the exposure was subtracted from the total volume inhaled. The difference indicates the amount by which the lungs were more filled, or less filled, at the end of the exposure than at the beginning. The differences were not greater

CONFIDENTIAL

CONFIDENTIAL

in either direction than the volume of a large breath and so may be accounted for by the possibility that the respiratory cycle of inspiration and expiration was not completed during the exposure. Minute volumes based on complete cycles were calculated for the periods before and during exposure. They were less during than before exposure but all were greater than the expected value for a man standing at ease. The maximum instantaneous inspiratory flow rate was recorded for each exposure and ranged from 50 liters per minute for relatively quiet breathing in BM to 250 liters per minute in LT. The number of coughs were counted during exposure and recovery; the triple exhalation in figure 7 (top record) was counted as one cough. Although coughing and dose were not systematically related, it is worth noting that LT and S had both doses and numbers of coughs that were far greater than those seen in the other exposures.

(U) Table 2 also includes the case of JC who was provided with a gas mask and instructed to put it on when he became aware of the CS. His breathing record during exposure is shown in figure 8, appendix. The cloud arrived at A during a normal exhalation of 1.77 liters. After inhaling only 0.17 liter the subject halted his breathing, opened the mask carrier at M and proceeded to don the mask. At C he exhaled 0.52 liter to clear the mask and resumed breathing. He was unable to breathe normally and after inhaling a further 2.07 liters he left the wind tunnel at B.

IV. (U) DISCUSSION.

In the large group of subjects described by Gutentag, Hart, Owens and Puntel from which our six subjects were drawn, the first symptom of CS exposure related to the respiratory tract was "burning" which began in the throat and progressed downward with or without coughing. As the exposure continued, this burning became painful and was rapidly followed by a "constricting" sensation throughout the chest. This latter symptom was associated with incapacitation for several minutes. Panic usually accompanied and accentuated this symptom and persons so affected appeared unable to inhale or exhale. Usually there was mild to moderate burning of the nose and throat. This was more irritating than incapacitating and disappeared readily in fresh air.

In the normal pneumotachograms of Fleisch,⁹ inspiration and expiration follow each other without an intervening period of zero flow. In a few normal subjects, Proctor and Hardy¹⁰ observed a pause of from 0.1 to 0.4 second following expiration. In our records the longest period of apnea was 4 seconds except for the longer voluntary breathhold of JC included in the masking reaction. Shorter pauses between the phases of

CONFIDENTIAL

UNCLASSIFIED

the respiratory cycle were quite common, however. Although there may have been some constriction of the airway, there was no reason to suppose the airway was blocked except momentarily as a part of the cough reflex.¹¹ The peak instantaneous flows in the coughs recorded here were not as high as the flows of from 500 to 700 liters per minute recorded by Whittenberger and Mead,¹² in part because of the limited range of the transducer.

The textbook description of the respiratory response to irritants is exemplified by the following statement of Schmidt:¹³ "A sneeze is elicited by irritation of the mucous membrane of the nose, a cough by irritation of the respiratory passages beyond the nose." He goes on to state that during anesthesia, irritation of the upper air passages causes apnea. This reflex is actually present in the conscious subject but is overshadowed by the violent expiratory effort that is absent in the unconscious state.

According to literature cited in reviews by Ranson¹⁴ and Aviado and Schmidt,¹⁵ the evidence regarding the sneeze is derived from experiments on anesthetized animals, usually the rabbit. Search of the literature has so far revealed no pneumotachogram of a human sneeze. In our records we saw nothing that looked like a frank sneeze, which we suppose would be characterized by a large slow inspiration followed by a sharp expiration, probably not as explosive as a cough. Sneezing was common, however, among observers exposed to small concentrations of CS at some distance from the exposure chamber. The impression of the pauses between inspiration and expiration in subject FC is that they were largely involuntary. The gasping behavior subjectively resembled what happens before a sneeze, but the irritation was more pronounced than that usually leading to a sneeze.

Electrical stimulation of the tracheobronchial mucosa accomplished during bronchoscopy indicates that the linings of the trachea and bronchi are sensitive and that pain results. Morton et al.¹⁶ have charted the areas in which the pain was felt. Pain arising from the bronchial tree was referred to the homolateral anterior chest within 2 to 4 centimeters of the sternum, or to the anterior cervical region within 2 centimeters of the midline. Pain of tracheal origin was felt in the midline anteriorly, at sites extending from the larynx to the xyphoid process. Pain of bronchial origin was usually abolished by a unilateral vagus section on the same side,¹⁷ and the cough reflex elicited by mechanical and electrical stimulation on the same side of the bronchial tree was also abolished.¹⁸

For the particle size (3 microns) employed on the 4 days of these tests, data cited by Punte¹⁹ indicate that as much as three-fourths of the aerosol would be retained in the upper respiratory tract. For particle sizes

UNCLASSIFIED

CONFIDENTIAL

of 1 micron and below, virtually all the material retained would be in the lower part of the tract. The smaller particle size was employed in some of the tests with the larger group of subjects.¹ It appears that either pain or coughing may result from stimulation of the tracheobronchial mucosa with either CS aerosol or an electric shock but we do not know the parameters of the stimulus that determine which result will occur. In view of the distribution of the particle size in these exposures, we cannot exclude the likelihood of penetration of some CS into the lower respiratory tract and may with Punte (personal communication) attribute to this the burning and constricting sensation of the chest. On the basis of the doses in table 2, it appears likely that a massive penetration is required to produce a bout of coughing. The statement of Schmidt,¹³ quoted above, suggests that the brief periods of apnea in our records result from strong stimulation in the nose. Much weaker stimulation with CS may be required to produce a sneeze.

Although CS produced great distortion of the pneumotachogram, the data for minute volume in table 2 make it clear that adequate ventilation of the lungs was maintained during these exposures. The large minute volume before exposure is interpreted as emotional hyperventilation.

The behavior of the man who donned the protective mask after the cloud reached him needs to be considered, for the respiratory flowmeter and its connecting cable could easily impair the seal of a hastily donned mask. However, the exposure time of JC was within the range of the untrained group provided with masks who were exposed to about the same concentration, described by Gutentag, Hart, Owens and Punte.¹ In a personal communication, Mr. A. L. West stated that a large breath was required to clear the mask after donning in order to render tolerable a long period of exposure to CS. If a good seal to the face could be obtained over the flowmeter, some quantitative studies on this point might be appropriate.

V. (C) SUMMARY.

1. (C) At a concentration of 5 micrograms per liter one trained subject was able to maintain a normal breathing pattern for 110 seconds.
2. (U) At higher concentrations the breathing pattern consisted of short inspiratory or expiratory gasps separated by short and variable periods of zero air flow.
3. (U) Coughing was conspicuous in only two exposures with the highest total inhaled doses.

CONFIDENTIAL

UNCLASSIFIED

4. (U) Ventilation of the lungs was maintained during the exposures at a normal or greater than normal rate.

5. (U) Voluntary Breath holding was not employed as a means of prolonging the exposure.

VI. (U) CONCLUSION.

Although the breathing patterns were disrupted by CS, adequate ventilation of the lungs was maintained, so that the incapacitation is attributed to the unpleasant sensations rather than to any degree of respiratory failure.

VII. (U) LITERATURE CITED.

1. Gutentag, P. J., Hart, J., Owens, E. J., and Punte, C. L. CWLR 2365. Evaluation of CS Aerosols as a Riot Control Agent in Man. April 1960. Confidential Report.

2. Dahlke, H. E., and Welkowitz, W. Diagnosing with Strain Gages. Electronic Industries 18, No. 1, 74-78 (1959). Unclassified Report.

3. Welkowitz, W., and Dahlke, H. Final Report. Contract DA-18-108-CML-5948, Gulton Industries, Inc., Metuchen, N. J. Portable Respiratory Volume Recording System. April 1957. Unclassified Report. See also subsequent reports by the same company on contract DA18-108-405-CML-52.

4. Cummings, E. G., Craig, F. N., and Blevins, W. V. CWLR 2323. Physiological Evaluation of the E52R26 Civilian Protective Mask. November 1959. Unclassified Report.

5. Upham, J. L., and Dranetz, A. I. Transistor Modulator for Airborne Recording. Electronics. pp.166-169. (June 1956). Unclassified Report.

6. Traite, M. Eighth Bi-Monthly Report. Contract DA-18-108-405-CML-52, Gulton Industries, Inc., Metuchen, N. J. January 1959. Unclassified Report.

7. Cummings, E. G., and Craig, F. N. Physiological Assessment, Appendix D in CWLR 2183. Mask, Protective, Field, E13R9, by I. S. Sherman. December 1957. Confidential Report.

UNCLASSIFIED

UNCLASSIFIED

8. Cummings, E. G., Blevins, W. V., and Craig, F. N. Measurement of an External Dead Space with a New Flowmeter. J. Appl. Physiol. 15, (1960). Unclassified Report.
9. Fleisch, A. Der Pneumotachograph; ein Apparat zur Geschwindigkeitsregistrierung der Atemluft. Pflüger's Arch. ges. Physiol. 209, 713-722 (1925). Unclassified Report.
10. Proctor, D. F., and Hardy, J. B. Studies of Respiratory Air Flow. I. Significance of the Normal Pneumotachogram. Bull. Johns Hopkins Hosp. 85, 258-280 (1949). Unclassified Report.
11. Dayman, H. Mechanics of Airflow in Health and in Emphysema. J. Clin. Invest. 30, 1175-1190 (1951). Unclassified Report.
12. Whittenberger, J. L., and Mead, J. Respiratory Dynamics During Cough. Trans. Natl. Tuberc. Assoc. 48, 414-418 (1952). Unclassified Report.
13. Schmidt, C. F. p. 380, in Bard's Medical Physiology. 10th Ed. St. Louis. 1956. Unclassified Report.
14. Ranson, S. W. Afferent Paths for Visceral Reflexes. Physiol. Rev. 1, 477-522 (1921). Unclassified Report.
15. Aviado, D. M., Jr., and Schmidt, C. F. Reflexes from Stretch Receptors in Blood Vessels, Heart and Lungs. Physiol. Rev. 35, 247-300 (1955). Unclassified Report.
16. Morton, D. R., Klassen, K. P., and Curtis, G. M. The Clinical Physiology of the Human Bronchi. I. Pain of Tracheobronchial Origin. Surgery 28, 699-704 (1950). Unclassified Report.
17. Morton, D. R., Klassen, K. P., and Curtis, G. M. The Clinical Physiology of the Human Bronchi. II. The Effect of Vagus Section upon Pain of Tracheobronchial Origin. Surgery 30, 800-809 (1951). Unclassified Report.
18. Klassen, K. P., Morton, D. R., and Curtis, G. M. The Clinical Physiology of the Human Bronchi. III. The Effect of Vagus Section on the Cough Reflex, Bronchial Caliber and Clearance of Bronchial Secretions. Surgery 29, 483-490 (1951). Unclassified Report.

UNCLASSIFIED

UNCLASSIFIED

19. Punte, C. L. Some Aspects of Particle Size in Aerosol Studies. Armed Forces Chem. J. March-April, 1958. Unclassified Report.

UNCLASSIFIED

CONFIDENTIAL

(C)

APPENDIX

TABLES AND GRAPHS

(U)

TABLE 1

COMPARISON OF TIDAL VOLUMES RECORDED WITH A SPIROMETER AND WITH THE PORTABLE TAPE RECORDING SYSTEM* (U)

Breath	Condition	Time 1330 mm min	Inspired volume		Expired volume
			Spirometer liters	Tape liters	Tape liters
1	Run	2			1.19
1	Run	42	1.19	1.48	
2	Run	85			1.27
2	Run	130	1.46	1.71	
3	Run	178			1.44
3	Run	228	1.46	1.69	
4	Run	274			1.44
4	Run	321	1.33	1.68	
5	Run	363			1.31
5	Run	404	1.19	1.26	
6	Stand	440			1.03
6	Stand	469	2.52	2.83	
7	Stand	535			2.32
7	Stand	598	1.59	1.69	
8	Stand	660			
Total			10.74	12.34	10.00

* Data were obtained in connection with another series of experiments on 3 March 1959 from subject F running at 9 miles per hour up a 12 per cent grade and include the first two breaths in recovery. (Craig, F. N., and Cummings, E. G. Breathing in Brief Exercise. 1960 manuscript).

CONFIDENTIAL

(C)

TABLE 2

SUMMARY OF EXPOSURE TO CS (U)

Date, 1960	4/3	3/30	4/6	3/30	4/6	4/3	3/31
Subject	BM	FC	LT	FC	K	S	JC**
Previous exposures	+	-	+	+	-	+	-
Concentration of CS, micrograms per liter*	5	12	15	64	80	150	175
Duration of exposure, seconds	110	24	61	15	12	12	27
Total dose inhaled, micrograms	101	55	451	129	184	450	-
Respiratory response time, seconds	-	5.2	11.6	4.2	4.6	4.9	2.6
Volume inhaled before response, liters	-	1.4	13.3	1.1	1.6	3.0	0.2
Time without breathing, seconds	0	9.2	12.0	5.8	8.2	4.5	19.1
Total volume inspired, liters	20.1	4.6	30.1	2.0	2.3	3.0	2.3
Inspiration less expiration, liters	-	1.1	1.9	-1.3	1.5	-2.2	0.7
Maximum inspiratory flow, liters per minute	50	95	250	200	70	160	150
Minute volume before exposure, liters per minute	21.7	13.1	23.5	17.4	20.7	34.4	17.3
Minute volume during exposure, liters per minute	11.0	16.5	29.6	8.0	11.5	15.0	5.1
Number of coughs	0	4	15	2	0	16	0

* The mass median diameter in these exposures was 3 microns.

** JC donned a protective mask when the cloud arrived.

CONFIDENTIAL

CONFIDENTIAL

UNCLASSIFIED




FIGURE 1

RECORDING AND ELECTRONICS SECTION OF PORTABLE BREATH-
RECORDING SYSTEM WITH COVER REMOVED TO
EXPOSE THE TAPE-TRANSPORT MECHANISM

UNCLASSIFIED

UNCLASSIFIED



FIGURE 2

PORTABLE BREATH-RECORDING SYSTEM WITH TRANSDUCER IN
POSITION ON THE FACE, AND KNAPSACK CONTAINING
THE POWER SUPPLY, REMOTE CONTROL RADIO
RECEIVER AND RECORDER SECTIONS

The M9A1 protective mask and carrier are wired to the remote control radio receiver section so that the times of opening the carrier and closing the neck strap of the mask can be noted on the record. The remote control radio transmitter shown at the left is used to start the tape recorder and to place an additional time signal on the tape.

UNCLASSIFIED

UNCLASSIFIED




FIGURE 3

CALIBRATION AND RESISTANCE OF VANE-TYPE STRAIN-GAGE AIR-FLOW TRANSDUCER

Electrical output is shown in pen deflection on the oscillograph as generated by the Sanborn strain-gage amplifier (not the tape recorder system), with an arbitrary scale setting of 25 millimeters for a flow of 250 liters per minute. Resistance for each rate of flow is given in millimeters of water pressure.

UNCLASSIFIED

UNCLASSIFIED




FIGURE 4

Complete records of the first three exposures listed in table 2 are shown. The time scale at the bottom of each record is in intervals of 2 seconds. Zero air flow is three lines down from the top of the record. Inspiration is down and expiration up from this line. Each interval between lines on the vertical scale represents an instantaneous air flow of 50 liters per minute. The subject entered the wind tunnel at the beginning of the record. At A the cloud arrived and at B the subject left the wind tunnel.

UNCLASSIFIED

UNCLASSIFIED



FIGURE 5

Shown are expanded portions of the records of figure 4 in the same order. The time scale at the bottom of each record is in intervals of 2 seconds. The scale for instantaneous air flow is the same as in figure 4. The remote control signal for the arrival of the cloud is seen at A.

UNCLASSIFIED

UNCLASSIFIED




FIGURE 6

Complete records of the second three exposures listed in table 2 are shown. Scales and symbols are as shown in figure 4.

UNCLASSIFIED

UNCLASSIFIED




FIGURE 7

Expanded portions of the records of figure 6 are shown. Scales and symbols are as shown in figure 5.

UNCLASSIFIED

UNCLASSIFIED




FIGURE 8

Shown are complete and expanded portions of the record of the last exposure in table 2 (JC). Scales and symbols are as shown in figures 4 and 5. The signal at M came from the gas mask carrier; when the carrier was opened the circuit was broken. C marks an exhalation to clear the mask after donning.

UNCLASSIFIED

UNCLASSIFIED

UNCLASSIFIED